



## A semi-automated method for measuring the evolution of both lumen area and blood flow in carotid from Phase Contrast MRI

Submitted by Jean-Baptiste F... on Mon, 06/18/2018 - 09:34

Titre	A semi-automated method for measuring the evolution of both lumen area and blood flow in carotid from Phase Contrast MRI
Type de publication	Article de revue
Auteur	Fasquel, Jean-Baptiste [1], Lécluse, Aldéric [2], Cavarro-Ménard, Christine [3], Willoteaux, Serge [4]
Editeur	Elsevier
Type	Article scientifique dans une revue à comité de lecture
Année	2015
Langue	Anglais
Date	1er Nov. 2015
Pagination	269-277
Volume	66
Titre de la revue	Computers in Biology and Medicine
ISSN	0010-4825
Mots-clés	Magnetic Resonance Imaging [5], Phantom [6], r Blood flow [7], Region-based active contou [8], Semi-automated analysis [9]
Résumé en anglais	<p>Phase-Contrast (PC) velocimetry Magnetic Resonance Imaging (MRI) is a useful modality to explore cardiovascular pathologies, but requires the automatic segmentation of vessels and the measurement of both lumen area and blood flow evolutions. In this paper, we propose a semi-automated method for extracting lumen boundaries of the carotid artery and compute both lumen area and blood flow evolutions over the cardiac cycle. This method uses narrow band region-based active contours in order to correctly capture the lumen boundary without being corrupted by surrounding structures. This approach is compared to traditional edge-based active contours, considered in related works, which significantly underestimate lumen area and blood flow. Experiments are performed using both a sequence of a homemade phantom and sequences of 20 real carotids, including a comparison with manual segmentation performed by a radiologist expert. Results obtained on the phantom sequence show that the edge-based approach leads to an underestimate of carotid lumen area and related flows of respectively 18.68% and 4.95%. This appears significantly larger than weak errors obtained using the region-based approach (respectively 2.73% and 1.23%). Benefits appear even better on the real sequences. The edge-based approach leads to underestimates of 40.88% for areas and 13.39% for blood flows, compared to limited errors of 7.41% and 4.6% with our method. Experiments also illustrate the high variability and therefore the lack of reliability of manual segmentation.</p>
URL de la notice	<a href="http://okina.univ-angers.fr/publications/ua17086">http://okina.univ-angers.fr/publications/ua17086</a> [10]

DOI 10.1016/j.compbio.2015.09.017 [11]

Lien vers le document <https://www.sciencedirect.com/science/article/pii/S0010482515003273?via%...> [12]

---

## Liens

- [1] <http://okina.univ-angers.fr/j.fasquel/publications>
- [2] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=28440>
- [3] <http://okina.univ-angers.fr/c.menard/publications>
- [4] <http://okina.univ-angers.fr/s.willoteaux/publications>
- [5] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=6040>
- [6] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=24745>
- [7] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=24744>
- [8] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=24743>
- [9] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=24742>
- [10] <http://okina.univ-angers.fr/publications/ua17086>
- [11] <http://dx.doi.org/10.1016/j.compbio.2015.09.017>
- [12] <https://www.sciencedirect.com/science/article/pii/S0010482515003273?via%3Dihub>

Publié sur *Okina* (<http://okina.univ-angers.fr>)